arXiv:1702.01883v1 [math.NT] 7 Feb 2017

Irreducible characters with bounded root Artin conductor

Amalia Pizarro-Madariaga
Instituto de Matemáticas
Universidad de Valparaíso, Chile
amalia.pizarro@uv.cl

Abstract

In this work, we prove that the growth of the Artin conductor is at most, exponential in the degree of the character.

1 Introduction

Let K be an algebraic number field such that $K/\mathbb{Q}$ is Galois and let χ be the character of a linear representation of $\mathrm{Gal}(K/\mathbb{Q})$. We denote by f_χ the Artin conductor of χ. In [7], Odlyzko found lower bounds for f_χ by applying analytic methods to the Artin L-function. We have improved Odlyzko's lower bounds in [5] by using explicit formulas for Artin L-functions. In particular, if χ is an irreducible character of $\mathrm{Gal}(K/\mathbb{Q})$ and assuming that $\chi\overline{\chi}$ satisfies the Artin conjecture we obtained

$$f_\chi^{1/\chi(1)} \geq 4.73(1.648)^{\frac{(a_\chi - b_\chi)^2}{\chi(1)^2}} e^{-(13.34/\chi(1))^2},$$

where a_χ and b_χ are integers depending on χ.

It is important to assume the Artin Conjecture in order to control the poles of Artin L-functions. Furthermore, this bound is even better when we assume that $L(s, \chi\overline{\chi})$ satisfies the Generalized Riemann Hypothesis.

In this work, we study the growth of the root Artin conductor given by $f_\chi^{1/\chi(1)}$. This fact has been studied for the discriminant of a number field. If $n_0 = r_1 + 2r_2$, let d_n be the minimal discriminant of the fields K with degree n multiple of n_0 and such that $r_1(K)$ and $r_2(K)$ are in the same ratio as r_1, r_2. Let $\alpha(r_1, r_2) = \lim \inf_{n\to\infty} d_n^{1/n}$. Martinet considered number fields with infinite 2-class field towers and proved that [6]

$$\alpha(0, 1) < 93 \text{ and } \alpha(1, 0) < 1059.$$

We follow this idea and consider a number field K with infinite p-class field tower for some prime p. Under some technical conditions over K, we find an upper bound (just depending on K) for the root Artin conductor of the irreducible characters of $\mathrm{Gal}(K_n/\mathbb{Q})$, where K_n is the Hilbert p-class field of K_{n-1} with $K_0 = K$.

More precisely, we propose in §2 a technique which is useful to classify the irreducible characters of $\mathrm{Gal}(K_n/\mathbb{Q})$ in terms of a certain normal subgroup. This characterization is convenient to obtain upper bounds for root Artin conductors.

We prove in §3 that for each $n \geq 1$ there exist irreducible characters χ of $\mathrm{Gal}(K_n/\mathbb{Q})$ whose degree $\chi(1)$ increase with n and such that

$$f_\chi^{1/\chi(1)} \leq C,$$

where $C > 0$ is an effective computable constant depending only of the number field K.

In §4 we apply the results obtained in §2 and §3 to the number field $K = \mathbb{Q}(\zeta_{11} + \zeta_{11}^{-1}, \sqrt{2}, \sqrt{-23})$. This field was found by Martinet in [6] and has infinite 2-class field tower and lowest known discriminant. In particular, we prove that for each $n \geq 1$ it is possible to find an irreducible character of $\mathrm{Gal}(K_n/\mathbb{Q})$ with large degree and

$$f_\chi^{1/\chi(1)} \leq C, \text{ where } C \leq 11^4 \cdot 2^{15} \cdot 23.$$

2 Irreducible characters with large degree

In this section, we develop a technique to classify the irreducible characters of groups with a normal subgroup of prime index. Also, we find conditions that ensure the existence of irreducible characters of large degree. We believe these results are of independent interest.

2.1 Clifford Theory

Let us consider a finite group G with a normal subgroup H such that $[G : H] = q$, where q is a prime number. We denote the set of the irreducible characters of G by $Irr(G)$. If χ and θ are characters of G and H respectively, we denote the restriction of χ to H by $Res_H^G\chi$ and the induced character of θ to G by $Ind_H^G\theta$. If $\theta \in Irr(H)$, we define the conjugate character to θ in G by $\theta^g : H \to \mathbb{C}$, where $\theta^g(h) = \theta(ghg^{-1})$. The inertia group of $\theta \in G$ is given by

$$I_G(\theta) = \{g \in G : \theta^g = \theta\}.$$

G acts on $Irr(H)$ by conjugation and $I_G(\theta)$ is the stabilizer of θ under this action.

Theorem 2.1. *(Clifford) Let H be a normal subgroup of G and $\theta \in Irr(H), \chi \in Irr(G)$ such that θ is an irreducible constituent of $Res_H^G\chi$, with $\langle Res_H^G\chi, \theta\rangle = e > 0$. Suppose that $\theta = \theta^{g_1}, \theta^{g_1}, \ldots, \theta^{g_j}$ are the distinct conjugates of θ in G. Assume also that*

$$G = \bigcup_{j=1}^{t} I_g(\theta)g_j, \text{ with } t = [G : I_G(\theta)].$$

Then,

(a) $Res_H^G Ind_H^G \theta = |I_G(\theta)/H| \sum_{j=1}^{t} \theta^{g_j}$.

(b) $\langle Ind_H^G \theta, Ind_H^G \theta \rangle = |I_G(\theta)/H|$. *In particular,* $Ind_H^G \theta \in Irr(G)$ *if and only if* $I_G(\theta) = H$.

(c) $Res_H^G \chi = e \sum_{j=1}^{t} \theta^{g_j}$. *In particular,*

$$\chi(1) = et\theta(1) \text{ and } \langle Res_H^G \chi, Res_H^G \chi \rangle = e^2 t.$$

Also, $e^2 \leq |I_G(\theta)/H|$ *and* $e^2 t \leq |G/H|$.

For details of the proof see [3],pp. 253.

We say that an irreducible character θ of H *is extendible to* G if there is an irreducible character χ of G such that $Res_H^G \chi = \theta$.

The following result of Gallagher [1], pp. 225 gives us a criterion to decide when a character is extendible.

Theorem 2.2. *Let* H *be a normal subgroup of prime index* q *in* G. *If* $\theta \in Irr(H)$ *is invariant in* G *(i.e.* $I_G(\theta) = G$*), then* θ *is extendible to* G.

2.2 Irreducible characters of G

Lemma 1. *Suposse that* G *has a normal subgroup* H *such that* $[G : H] = q$*, where* q *is a prime number. If* $\theta \in Irr(H)$*, then the inertia group of* θ *is either,*

(i) $I_G(\theta) = G$, *or*

(ii) $I_G(\theta) = H$.

Proof. Since H is a normal subgroup of G, $H \subset I_G(\theta)$. So if we put $t = [G : I_G(\theta)]$, then $t \mid [G : H]$. Therefore $t = 1$ or $t = q$. □

Theorem 2.3. *Under the conditions of Lemma 1, let* χ *be an irreducible character of* G. *Then either,*

(i) $Res_H^G \chi = \theta$, *for some* $\theta \in Irr(H)$, *or*

(ii) $\chi = Ind_H^G \theta$, *for some* $\theta \in Irr(H)$.

Proof. Let $\chi \in Irr(G)$ and take $\theta \in Irr(H)$ an irreducible constituent of $Res_H^G \chi$. Then $Res_H^G \chi = e \sum_{i=1}^{t} \theta_i$, where θ_i are the conjugates of θ and $\langle Res_H^G \chi, \theta \rangle = e > 0$. Let us fix $\theta = \theta_1$. If $t = 1$, then $I_G(\theta) = G$ and by the Theorem 2.2, θ is extendible to G. On the other hand we know that (see Theorem 19.4 in [3])

$$e^2 \mid [G : H] = q,$$

so $e^2 = 1$. Therefore, $e = 1$ and $Res_H^G \chi = \theta$, so we have (i).

If $t = q$, then $I_G(\theta) = H$ and from Theorem 2.1 $Ind_H^G \theta \in Irr(G)$. By Frobenius reciprocity,

$$\langle Ind_H^G \theta, \chi \rangle_G = \langle \theta, Res_H^G \chi \rangle_H = \overline{\langle Res_H^G \chi, \theta \rangle} = \overline{e} = e.$$

Since $Ind_H^G \theta$ and χ are irreducible and θ is a consituent of $Res_H^G \chi$, it follows that $e = 1$ and so $Ind_H^G \theta = \chi$.

□

2.3 Irreducible characters with large degree

In order to ensure the existence of irreducible characters of large degree, we prove the following lemma given as an exercise in [4], pp. 98.

Lemma 2. *Let G be a group with a chain of normal subgroups*

$$1 = H_0 \trianglelefteq H_1 \trianglelefteq H_2 \ldots \trianglelefteq H_n = G$$

such that H_i/H_{i-1} is non abelian for $i = 1, \ldots, n$. Then, there exists an irreducible character ϕ of G, such that $\phi(1) \geq 2^n$.

Proof. By induction over n. For $n = 1$, note that as $H_1 \cong H_1/H_0$ is non abelian there exists a character $\psi_1 \in Irr(H_1)$ with $\psi_1(1) \geq 2$.

For $n = 2$, let $\psi_2 \in Irr(H_2)$ such that ψ_1 is an irreducible constituent of $Res_{H_1}^{H_2} \psi_2$ (take for example $\psi_2 \in Ind_{H_1}^{H_2} \psi_1$). From Theorem 2.1, it follows

$$Res_{H_1}^{H_2} \psi_2 = e \sum_{i=1}^{t} \psi_{1,i}, \tag{2.1}$$

where $\psi_{1,1} = \psi_1, \ldots, \psi_{1,t}$ are the distinct conjugates of ψ_1 in H_2, $e = \langle Res_{H_1}^{H_2} \psi_2, \psi_1 \rangle$ and $t = [H_2 : I_{H_2}(\psi_1)]$.

If $t = 1$, then $H_2 = I_{H_2}(\psi_1)$ and $Res_{H_1}^{H_2} \psi_2 = e\psi_1$. Hence,

(a) $\psi_2(1) = e\psi_1(1) \geq 2$.

(b) For all $\beta \in Irr(H_2/H_1)$ the character $\psi_2\beta$ belongs to $Irr(H_2)$ (see Theorem 19.5 in [3]). As H_2/H_1 is non abelian, we choose $\beta_2 \in Irr(H_2/H_1)$ such that $\beta_2(1) \geq 2$. Then $\phi = \psi_2\beta_2 \in Irr(H_2)$ and $\phi(1) = \psi_2(1)\beta_2(1) \geq 2^2$.

If $t \geq 2$, we just take $\phi = \psi_2$ so, $\phi(1) = et\psi_1(1) \geq 2\psi_1(1) \geq 2^2$.

Now, suppose that it is true for every $m < n$. Then there is $\psi_m \in Irr(H_m)$ such that $\psi_m(1) \geq 2^m$. Let us choose $\psi_{m+1} \in Irr(H_{m+1})$ such that ψ_m is an irreducible constituent of $Res_{H_m}^{H_{m+1}} \psi_{m+1}$. As in (2.1)

$$Res_{H_m}^{H_{m+1}} \psi_{m+1} = e \sum_{i=1}^{t} \psi_{m,i},$$

where $\psi_{m,1} = \psi_m, \ldots, \psi_{m,t}$ are the conjugates of ψ_m in H_{m+1}, $e = \langle Res^{H_m}_{H_{m+1}} \psi_{m+1}, \psi_m \rangle$ and $t = [H_{m+1} : I_{H_{m+1}}(\psi_m)]$.

If $t = 1$, then

(a) $\psi_{m+1}(1) \geq 2^m$;

(b) $\psi_{m+1}\beta \in Irr(H_{m+1})$, with $\beta \in Irr(H_{m+1}/H_m)$. Let us choose β such that $\beta(1) \geq 2$, then $\phi = \beta\psi_{m+1} \in Irr(H_{m+1})$ and $\phi(1) = \beta(1)\psi_{m+1}(1) \geq 2^{m+1}$.

If $t \geq 2$, we take $\phi = \psi_{m+1}$ and $\phi(1) \geq et\psi_m(1) \geq 2^{m+1}$.

□

Lemma 3. *Let H be a normal subgroup of G. If H has an irreducible character θ with $\theta(1) \geq r$, then there exists $\chi \in Irr(G)$ with $\chi(1) \geq r$.*

Proof. Let $\theta \in Irr(H)$ with $\theta(1) \geq r$ and choose $\chi \in Irr(G)$ such that θ is an irreducible constituent of $Res^G_H \chi$. By Theorem 2.1, $\chi(1) = et\theta(1)$, where $e = \langle Res^G_H \chi, \theta \rangle$ and $t = [G : I_g(\theta)]$. As $e, t \geq 1$, then $\chi(1) \geq \theta(1) \geq r$. □

3 Estimation for the root Artin conductor of irreducible characters of G_n

In this section, we aim to prove that the growth of the Artin conductor is at most, exponential in the degree of the character. It is well known that if L is a Galois unramified extension of K and ρ is a linear representation of $\mathrm{Gal}(L/M)$, then f_ρ is a trivial ideal. Then, in order to find a family of irreducible representations with bounded root Artin conductor it is convenient to analyse number fields with infinite p-class field tower, for some prime p.

Let us suposse K is a number field with infinite p-class field tower, K_n the Hilbert p-class field of K_{n-1} with $K_0 = K$ and G_n=$\mathrm{Gal}(K_n/\mathbb{Q})$. Under some conditions over K and applying the results of the previous section, we obtain that for each $n > 1$ it is possible to find an irreducible character of G_n with bounded root Artin conductor and degree increasing with n.

Theorem 3.1. *Let K be a number field Galois over $\mathbb{Q}$ with infinite p-class field tower, for some prime p. Suppose that K has a subfield $\widetilde{k}$ satisfying the following conditions:*

(a) $\widetilde{k}$ is Galois over $\mathbb{Q}$.

(b) $[\widetilde{k} : \mathbb{Q}] = q$, with q a prime number.

Let $\chi \in Irr(G_n)$, where $G_n = Gal(K_n/\mathbb{Q})$. If $\widetilde{H}_n = Gal(K_n/\widetilde{k})$, then either

(i) $Res^{G_n}_{\widetilde{H}_n} \chi = \theta$, for some $\theta \in Irr(\widetilde{H}_n)$, or

(ii) $\chi = Ind^{G_n}_{\widetilde{H}_n} \theta$, for some $\theta \in Irr(\widetilde{H}_n)$.

Proof. The proof follows directly from the Theorem 2.3 with $G = G_n$ and $H = \widetilde{H}_n$. □

Proposition 1. *Assume G_n under conditions of Theorem 3.1. Then for each $n \geq 1$, there exits $\chi \in Irr(G_n)$ such that*

$$\chi(1) > 2^{\frac{n-1}{2}}.$$

Proof. We fix $n \geq 1$. By Lemma 3, it is enough to show that if $G = Gal(K_n/K)$, then there exists $\phi \in Irr(G)$ such that $\phi(1) \geq 2^{\frac{n-1}{2}}$. In fact, let us consider the following chain of subgroups. If n is even, we take for $1 \leq j \leq \frac{n}{2}$:

$$\begin{aligned}
H_0 &= \{1\}, \\
H_1 &= Gal(K_n/K_{n-2}), \quad H_1/H_0 \cong H_1 \\
H_2 &= Gal(K_n/K_{n-4}), \quad H_2/H_1 \cong Gal(K_{n-2}/K_{n-4}) \\
&\vdots \\
H_j &= Gal(K_n/K_{n-2j}), \quad H_j/H_{j-1} \cong Gal(K_{n-2(j-1)}/K_{n-2j}) \\
&\vdots \\
H_{\frac{n}{2}} &= G = Gal(K_n/K), \quad H_{\frac{n}{2}}/H_{\frac{n}{2}-1} \cong Gal(K_2/K).
\end{aligned}$$

If $l < i-1$ then K_i/K_l is a non abelian group so, by Lemma 2 there exists $\phi \in Irr(G)$ with $\phi(1) \geq 2^{\frac{n}{2}} > 2^{\frac{n-1}{2}}$. If n is odd, for $j < \frac{n-1}{2}$ we take H_j and H_j/H_{j-1} as in the even case. For $j = (n-1)/2$ we take $H_{\frac{n-1}{2}}$=G and $H_{\frac{n-1}{2}}/H_{\frac{n-1}{2}-1} \cong Gal(K_3/K)$. Hence, there exists $\phi \in Irr(G)$ such that $\phi(1) \geq 2^{\frac{n-1}{2}}$.

□

Let us recall the definition of the Artin conductor. If L/M is a Galois extension and χ is the character of a linear representation of $\mathrm{Gal}(L/M)$, the Artin conductor is given by the ideal

$$f_\chi = \prod_{\mathfrak{p} \nmid \infty} \mathfrak{p}^{f_\chi(\mathfrak{p})}$$

with

$$f_\mathfrak{p}(\chi) = \frac{1}{|G_0|} \sum_{j \geq 0} (|G_j|\chi(1) - \chi(G_j)),$$

where G_i is the i-th ramification group of the local extension $L_\mathfrak{b}/M_\mathfrak{p}$ with $\mathfrak{b}$ a prime over $\mathfrak{p}$ and $\chi(G_j) = \sum_{g \in G_j} \chi(g)$.

Now, we are able to obtain upper bounds for the root Artin conductor of irreducible characters of G_n.

Theorem 3.2. *Assume G_n under conditions of Theorem 3.1 and $\chi \in Irr(G_n)$.*

(i) If $Res^{G_n}_{\widetilde{H}_n} \chi = \theta$, for some $\theta \in Irr(\widetilde{H}_n)$ then

$$f_\chi^{1/\chi(1)} \leq |D_{\widetilde{k}/\mathbb{Q}}|^{1/q} N_{\widetilde{k}/\mathbb{Q}}(f_\theta)^{1/\theta(1)}.$$

(ii) If $\chi = Ind_{\widetilde{H}_n}^{G_n}\theta$, *for some* $\theta \in Irr(\widetilde{H}_n)$ *then*

$$f_\chi^{1/\chi(1)} = |D_{\widetilde{k}/\mathbb{Q}}|^{1/q} N_{\widetilde{k}/\mathbb{Q}}(f_\theta)^{1/q\theta(1)}.$$

Proof. In the first case, we have $\chi(1) = \theta(1)$ and

$$Ind_{\widetilde{H}_n}^{G_n}\theta = \sum_{i=1}^{q} \psi_i(1) \cdot \chi\psi_i,$$

where $Irr(G_n/\widetilde{H}_n) = \{\psi_1, \psi_2, \ldots, \psi_q\}$ (see Theorem 19.5 in [3]). Since $G_n/\widetilde{H}_n$ is isomorphic to the abelian group $\mathbb{Z}/q\mathbb{Z}$, it follows $Ind_{\widetilde{H}_n}^{G_n}\theta = \sum_{i=1}^{q} \chi\psi_i$. The Artin conductor of this induced character is, on the one hand

$$f_{Ind_{\widetilde{H}_n}^{G_n}\theta} = |D_{\widetilde{k}/\mathbb{Q}}|^{\theta(1)} N_{\widetilde{k}/\mathbb{Q}}(f_\theta),$$

where the ideal f_θ is the Artin conductor of θ. On the other hand, assuming that ψ_1 is the trivial character

$$f_{Ind_{\widetilde{H}_n}^{G_n}\theta} = f_{\sum_{i=1}^{q}\chi\psi_i} = f_\chi \cdot \prod_{i=2}^{q} f_{\chi\psi_i}.$$

Now, combining these expressions we get

$$f_\chi = |D_{\widetilde{k}/\mathbb{Q}}|^{\theta(1)} N_{\widetilde{k}/\mathbb{Q}}(f_\theta) \cdot \left(\prod_{i=2}^{q} f_{\chi\psi_i}\right)^{-1},$$

so

$$f_\chi^{1/\chi(1)} \leq |D_{\widetilde{k}/\mathbb{Q}}| N_{\widetilde{k}/\mathbb{Q}}(f_\theta)^{1/\theta(1)}.$$

In the second case,

$$\chi(1) = [G_n : \widetilde{H}_n]\theta(1) = q\theta(1)$$

and we can see that the root Artin conductor of χ is given by the expression

$$f_\chi^{1/\chi(1)} = |D_{\widetilde{k}/\mathbb{Q}}|^{1/q} N_{\widetilde{k}/\mathbb{Q}}(f_\theta)^{1/q\theta(1)}.$$

□

Corollary 1. *For each* $n \geq 1$, *there exists an irreducible character* χ *of* G_n *with degree greater than* $2^{\frac{n-1}{2}}$ *such that*

$$f_\chi^{1/\chi(1)} \leq C,$$

where $C = C(\widetilde{k}) > 0$ *is an effective computable constant.*

Proof. By the Proposition 1 and the previous Theorem, we know that for each n there is an irreducible character χ of G_n with large degree and

$$f_\chi^{1/\chi(1)} \leq |D_{\widetilde{k}/\mathbb{Q}}| N_{\widetilde{k}/\mathbb{Q}}(f_\theta)^{1/\theta(1)},$$

for some $\theta \in Irr(\widetilde{H}_n)$. From properties of the higher ramification groups, it is possible to prove that $N_{\widetilde{k}/\mathbb{Q}}(\mathfrak{p}) \leq T^{\theta(1)}$, for some $T > 0$ depending only on the field $\widetilde{k}$. Hence,

$$f_\chi^{1/\chi(1)} \leq |D_{\widetilde{k}/\mathbb{Q}}| T := C.$$

□

4 Number fields with infinite 2-class field tower

Golod and Shafarevich [2] proved that a number field K has a infinite p-class field tower if the p-rank of the class group of K is large enough. In this case

$$\alpha(r_1, r_2) \leq |D_K|^{1/[K:\mathbb{Q}]},$$

where D_K is the discriminant of K.

In addition, Martinet has constructed a number field with infinite Hilbert class field towers and lowest known root discriminant and proved that

$$\alpha(0,1) < 93 \text{ and } \alpha(1,0) < 1059.$$

In particular, he found that $K = \mathbb{Q}(\zeta_{11} + \zeta_{11}^{-1}, \sqrt{2}, \sqrt{-23})$ has infinite 2-class field tower. Since $\widetilde{k} = \mathbb{Q}(\zeta_{11} + \zeta_{11}^{-1})$ is a subfield of K of degree 5 over $\mathbb{Q}$, K satisfies the conditions of the Theorem 3.1. The discriminant of $\widetilde{k}$ is

$$|D_{\widetilde{k}/\mathbb{Q}}| = 14641 = 11^4$$

and the only rational primes that ramify in K are 2, 11 and 23. Using PARI/GP [8], we can estimates the sizes of the higher ramification groups. We get the upper bound

$$N_{\widetilde{k}/\mathbb{Q}}(f_\theta) \leq (2^{15}23)^{\theta(1)}.$$

With this estimation it follows the explicit result:

Corollary 2. *For each $n \geq 1$, there exists a irreducible character χ such that $\chi(1)$ increase with n and*

$$f_\chi^{1/\chi(1)} \leq C, \text{ where } C \leq 11^4 \cdot 2^{15} \cdot 23.$$

An open problem now is to improve the constant C.

Acknowledgements. The author thanks to Eduardo Friedman for suggesting this problem and for helpful advice. The author also thanks Bill Allombert, Yuri Bilu, Mariela Carvacho and Andrea Vera for useful discussions. The author was partially supported by Red Iberoamericana de Teoría de Números and IdEx-Université de Bordeaux (credits of Benjamin Matschke). While working on this project, the author visited the ICMAT Madrid and IMB Bordeaux.

References

[1] P. X. Gallagher, *Group characters and normal Hall subgroups*, Nagoya Math. J. **21** (1962), 223–230.

[2] E. S. Golod and I. R. Šafarevič, *On the class field tower*, Izv. Akad. Nauk SSSR Ser. Mat. **28** (1964), 261–272.

[3] B. Huppert, *Character theory of finite groups*, De Gruyter, Berlin-New York, 1998.

[4] I. Martin Isaac, *Character theory of finite groups*, AMS Chelsea Publishing, 2006.

[5] A. Pizarro – Madariaga, *Lower bounds for the Artin conductor*, Mathematics of Computation **80** (2011), no. 273, 539–561.

[6] J. Martinet, *Tours de corps de classes et estimations de discriminants*, Inventiones Mathematicae **44** (1978), 65–73.

[7] A.M. Odlyzko, *On conductors and discriminants*, Proc. Sympos., Univ. Durham **1** (1977), 377–407.

[8] The PARI Group, Bordeaux, *PARI/GP version* 2.7.1, 2014, available from http://pari.math.u-bordeaux.fr/.